

q -Stokes Phenomenon of a Basic Hypergeometric Series ${}_1\phi_1(0; a; q, x)$

By

Yousuke OHYAMA

*Department of Mathematical Sciences
Tokushima University
Tokushima 770-8506, JAPAN
e-mail : ohyama@tokushima-u.ac.jp*

October 9, 2016

Abstract

We show a connection formula of a linear q -differential equation satisfied by ${}_1\phi_1(0; a; q, x)$. The basic hypergeometric series ${}_1\phi_1(0; a; q, x)$ represents the Hahn-Exton q -Bessel function. Since the q -differential equation has a divergent series solution, a q -analogue of the Stokes phenomenon appears. We give a resummation procedure of the divergent series by means of the q -Borel-Laplace transformation of order $1/2$.

2010 Mathematics Subject Classification. 34M40, 33D15.

Introduction

We study the following q -difference equation

$$ay(qz) + [z - (a + q)]y(z) + qy(z/q) = 0, \quad (1)$$

which has a solution $y(z) = {}_1\phi_1(0; a; q, z)$. We assume that $a \neq 0$. The basic hypergeometric series ${}_1\phi_1(0; a; q, z)$ is related to the Hahn-Exton q -Bessel function [15], one of the three different types of Jackson's q -analogue of the Bessel function. We solve the connection problem of (1), which gives relations between solutions around the origin and solutions around the infinity. Since (1) has a solution represented by a divergent power series around the infinity, a q -analogue of the Stokes phenomenon appears when we give a resummation of the divergent series. We show a resummation of the divergent solution by means of the q -Borel-Laplace transformation of order $1/2$, which is studied by Dreyfus and Eloy [1, 2].

It is known that there exist three different types of q -analogues of the Bessel function. Jackson defines his first q -analogue of the Bessel functions in [6], and the second q -Bessel function is introduced in [7]. Following the modern notation by Ismail [5], we denote

$$\begin{aligned} J_\nu^{(1)}(x; q) &= \frac{(q^{\nu+1}; q)_\infty}{(q; q)_\infty} \left(\frac{x}{2}\right)^\nu {}_2\phi_1\left(0, 0; q^{\nu+1}; q, -\frac{x^2}{4}\right), \\ J_\nu^{(2)}(x; q) &= \frac{(q^{\nu+1}; q)_\infty}{(q; q)_\infty} \left(\frac{x}{2}\right)^\nu {}_0\phi_1\left(-; q^{\nu+1}; q, -\frac{q^{\nu+1}x^2}{4}\right), \\ J_\nu^{(3)}(x; q) &= \frac{(q^{2\nu+2}; q^2)_\infty}{(q^2; q^2)_\infty} x^\nu {}_1\phi_1\left(0; q^{2\nu+2}; q^2, x^2q^2\right). \end{aligned}$$

Since the third one is found by Hahn [4] and Exton [3], it is called the Hahn-Exton q -Bessel function, which satisfies the q -difference equation

$$f(xq^2) + q^{-\nu}(x^2q^2 - 1 - q^{2\nu})f(xq) + f(x) = 0. \quad (2)$$

Any linear q -difference equation has two singular points, the origin and the infinity. Local solutions around each singular point are represented by a product of theta functions and a formal power series. A connection problem of a linear q -difference equation is to give a relation between the system of local solutions at the origin and the infinity. When the power series is divergent, the q -Stokes phenomenon appears by a resummation procedure.

A connection formula of the first Jackson q -Bessel function is shown by Zhang [19]. Since the second Jackson q -Bessel function is related to the first Jackson q -Bessel function

$$J_\nu^{(2)}(x; q) = (-x^2/4; q)_\infty \cdot J_\nu^{(1)}(x; q),$$

the connection formula of $J_\nu^{(2)}(x; q)$ follows from the connection formula of the first Jackson q -Bessel function.

But the connection problem for the third Jackson q -Bessel function has not been solved completely. Solutions of the Hahn-Bessel equation (2) has two independent solutions represented by $J_\nu^{(3)}(x; q)$ around the origin. One local solution around the infinity is represented by a convergent power series, and the other is represented by a divergent power series. The asymptotic behavior of $J_\nu^{(3)}(x; q)$ around the infinity is studied by Olde Daalhuis [13], but the connection problem is not treated. One connection formula only for the convergent series around the infinity has been shown by Morita [11]. But the q -Stokes phenomenon of the divergent power series solution is not studied.

In section two we show a q -difference equation satisfied by ${}_1\phi_1(0; a; q, x)$. In section three we review the q -Borel transformation and q -Laplace transformation. In section four we give a resummation of the divergent solution of (1). For divergent power series which satisfy q -difference equations, the q -Borel-Laplace

transformation is a powerful tool to give a q -summation procedure [14]. The Newton diagram of (1) has two segments at the infinity. The slopes of two segments are 1 and -1 . Since the difference of the two slopes are two, we need the q -Borel resummation of order $1/2$ [1, 2]. For q -difference linear equations, the Stokes region is not an angle domain, but an open dense set $\mathbb{C}^* \setminus \lambda q^{\mathbb{Z}}$ for $\lambda \in \mathbb{C}^*$ [18, 14]. By using the q -Borel-Laplace resummation method of order $1/2$, we show the q -Stokes phenomenon of the divergent series solution of (1) and the q -Stokes region is not outside of a q -spiral $\lambda q^{\mathbb{Z}}$ but outside of a \sqrt{q} -spiral $\lambda\sqrt{q}^{\mathbb{Z}}$. We remark that a Borel transformation for q -series is also studied by Jackson [8].

In section five we show a connection formula for the convergent solution of (1) around the infinity. This formula is essentially shown in [11]. Thus we obtain a complete connection formula of (1).

In the case $a = -q$, (1) reduces to the q -Airy equation studied by Hamamoto, Kajiwara and Witte [10]. The q -Stokes phenomenon of the q -Airy equation is also studied by Morita [12]. Our results contains the connection formula for the q -Airy equation.

The author gives his gratitude to Professor Changgui Zhang and Professor Jean-Pierre Ramis for fruitful discussions. Most of works has done during his stay at Institut de Mathématiques de Toulouse on September 2016. The author also expresses his appreciation for the kind hospitality to Professor Jacques Sauloy. This work is supported by JSPS KAKENHI Grant-in-Aid for Scientific Research (C) Number 6K05176.

1 Notations and Preliminary

In the following we assume that $q \in \mathbb{C}^*$ and $0 < |q| < 1$. For $n = 0, 1, 2, \dots$, we set the q -shifted factorial

$$(a; q)_n = \prod_{j=0}^{n-1} (1 - aq^j), \quad (a; q)_\infty = \prod_{j=0}^{\infty} (1 - aq^j).$$

We set $(a_1, a_2, \dots, a_m; q)_n = \prod_{j=1}^m (a_j; q)_n$ for $n = 0, 1, 2, \dots$ or $n = \infty$.

We set the theta function

$$\theta_q(x) := \theta(x) = \sum_{k \in \mathbb{Z}} q^{k(k-1)/2} x^k = (q, -x, -q/x; q)_\infty.$$

The theta function satisfies

$$\begin{aligned} \theta(q^k x) &= q^{-k(k-1)/2} x^{-k} \theta(x) \quad (k \in \mathbb{Z}), \\ x\theta(1/x) &= \theta(x), \quad \theta(1/x) = \theta(qx). \end{aligned}$$

It is easy to show the following lemma on relations between different bases q .

Lemma 1. *We have*

$$\begin{aligned}(x; q)_\infty &= (x; q^2)_\infty (xq; q^2)_\infty, \\ (x; q)_\infty (-x; q)_\infty &= (x^2; q^2)_\infty, \\ (q^2; q^2)_\infty \theta_q(x) &= (q; q^2)_\infty \theta_{q^2}(x) \theta_{q^2}(xq), \\ (-q; q)_\infty \theta_q(x) \theta_q(-x) &= (q; q)_\infty \theta_{q^2}(-x^2).\end{aligned}$$

1.1 Transformation of q -difference equation

The q -difference operator σ_q is given by $\sigma_q[f(t)] = f(tq)$. We use the following lemma frequently in this paper. The proof is evident.

Lemma 2. *We transform a second order q -difference equation*

$$[a(z)\sigma_q + b(z) + c(z)\sigma_q^{-1}]y(z) = 0.$$

(1) *We set $t = 1/z$ and $v(t) = y(1/t)$. Then $v(t)$ satisfies*

$$[c(1/t)\sigma_q + b(1/t) + a(1/t)\sigma_q^{-1}]v(t) = 0.$$

(2) *We set $y(z) = \theta(rz)y_1(z)$. Then $y_1(z)$ satisfies*

$$\left[\frac{a(z)}{rz}\sigma_q + b(z) + \frac{rzc(z)}{q}\sigma_q^{-1} \right] y_1(z) = 0.$$

(3) *We set $y(z) = (rz; q)_\infty y_2(z)$. Then $y_2(z)$ satisfies*

$$\left[\frac{a(z)}{1-rz}\sigma_q + b(z) + (1-rz/q)c(z)\sigma_q^{-1} \right] y_2(z) = 0.$$

1.2 Basic hypergeometric series

The basic hypergeometric series [9] is defined by

$$\begin{aligned}& {}_r\phi_s(a_1, \dots, a_r; b_1, \dots, b_s; q, x) \\ & := \sum_{n \geq 0} \frac{(a_1, \dots, a_r; q)_n}{(b_1, \dots, b_s; q)_n (q; q)_n} \left\{ (-1)^n q^{\frac{n(n-1)}{2}} \right\}^{1+s-r} x^n.\end{aligned}$$

Heine's basic hypergeometric series ${}_2\phi_1(a, b; c; q, z)$ satisfies the equation

$$[(c - abqz)\sigma_q^2 - (c + q - (a + b)qz)\sigma_q + q(1 - z)] {}_2\phi_1(a, b; c; q, z) = 0.$$

A connection formula of ${}_2\phi_1(a, b; c; q, z)$ is shown by Thomae [16] and Watson [17]:

$$\begin{aligned}{}_2\phi_1(a, b; c; q; x) &= \frac{(b, c/a; q)_\infty}{(c, b/a; q)_\infty} \frac{\theta(-ax)}{\theta(-x)} {}_2\phi_1(a, aq/c; aq/b; q, cq/abx) \\ &+ \frac{(a, c/b; q)_\infty}{(c, a/b; q)_\infty} \frac{\theta(-bx)}{\theta(-x)} {}_2\phi_1(b, bq/c; bq/a; q, cq/abx).\end{aligned}\quad (3)$$

It is known that there exist many relations between hypergeometric series. The relation

$${}_0\phi_1(-; a^2q; q^2, a^2qx^2) = (x; q)_\infty \cdot {}_2\phi_1(a, -a; a^2; q, x) \tag{4}$$

is shown in [19].

1.3 Formal q -Borel transformation

We review the q -Borel transformation and the q -Laplace transformation. See [14, 18, 20] for detail.

The q -Borel transformation $\mathcal{B}_q^\pm : \mathbb{C}[[t]] \rightarrow \mathbb{C}[[\tau]]$ is defined by

$$\mathcal{B}_q^\pm \left[\sum_{n=0}^\infty a_n t^n \right] := \sum_{n=0}^\infty a_n q^{\pm n(n-1)/2} \tau^n.$$

In usual we identify a germ of holomorphic functions at the origin $\mathcal{O}_{\mathbb{C},0}$ as a subset of $\mathbb{C}[[t]]$. As a linear operator on $\mathbb{C}[[t]]$, we have

$$\mathcal{B}_q^\pm (t^m \sigma_q^n f) = q^{\pm m(m-1)/2} \tau^m \sigma_q^{n \pm m} \mathcal{B}_q^\pm (f).$$

The q -Laplace transform of $\varphi(\tau)$ is given by the Jackson integral

$$L_{q;1}^{[\lambda]} \varphi(t) := \frac{1}{1-q} \int_0^{\lambda\infty} \frac{\varphi(\tau)}{\theta_q(\tau/x)} \frac{d_q \tau}{\tau} = \sum_{n \in \mathbb{Z}} \frac{\varphi(q^n \lambda)}{\theta_q(q^n \lambda/x)}.$$

When $f(t) \in \mathbb{C}[[t]]$ is a convergent power series,

$$L_{q;1}^{[\lambda]} \circ \mathcal{B}_q^+(f) = f.$$

In this sense, $L_{q;1}^{[\lambda]}$ is a formal inverse of \mathcal{B}_q^+ .

The following lemma is useful to calculate the q -Laplace transform. We can prove by direct calculations.

Lemma 3. 1) Assume that

$$\varphi(\xi) = \frac{\theta(a\xi)}{\theta(b\xi)} \sum_{m \geq 0} c_m \xi^{-m}.$$

Then

$$L_{q;1}^{[\lambda]} \varphi(x) := \frac{\theta(a\lambda)\theta(qax/b\lambda)}{\theta(b\lambda)\theta(qx/\lambda)} \sum_{m \geq 0} c_m q^{-m(m-1)/2} (b/aqx)^m.$$

In the case $a = b$, we obtain

$$L_{q;1}^{[\lambda]} \left[\sum_{m \geq 0} c_m \xi^{-m} \right] = \sum_{m \geq 0} c_m q^{-(-m)(-m-1)/2} x^{-m},$$

which gives a formal q -Borel transformation \mathcal{B}_q^- .

2) Assume that

$$\varphi(\xi) = \frac{\theta(a\xi)}{\theta(b_1\xi)\theta(b_2\xi)} \sum_{m \geq 0} c_m \xi^{-2m}.$$

Then

$$L_{q;1}^{[\lambda]} \varphi(x) := \frac{\theta_q(a\lambda)\theta_{q^2}(aq^2x/b_1b_2\lambda^2)}{\theta_q(b_1\lambda)\theta_q(b_2\lambda)\theta_q(qx/\lambda)} \sum_{m \geq 0} c_m q^{-m(m-1)} (b_1b_2/aq^2x)^m.$$

2 q -difference equation satisfied by ${}_1\phi_1(0; a; q, z)$

We assume $a \neq 0$. We consider the following q -difference equation

$$ay(qz) + [z - (a + q)]y(z) + qy(z/q) = 0, \quad (5)$$

which has a solution $y(z) = {}_1\phi_1(0; a; q, z)$. Since the degree of the coefficients of (1) is up to one, we study (1) instead of the Hahn-Exton equation (2).

We set $t = 1/z$ and $v(t) = y(1/t)$. Then $v(t)$ satisfies

$$qtv(tq) + [1 - (a + q)t]v(t) + atv(t/q) = 0. \quad (6)$$

In the following, we study a connection problem and the q -Stokes phenomenon of (6). Since (6) has a divergent series solution around the origin, the q -Stokes phenomenon appears when we give a resummation of the divergent series.

Local solutions of (6) around $t = \infty$ are

$$v_1^{(\infty)}(t) = {}_1\phi_1(0; a; q, 1/t), \quad v_2^{(\infty)}(t) = \frac{\theta(-qt)}{\theta(-at)} {}_1\phi_1(0; q^2/a; q, q/at).$$

Local (formal) solutions of (6) around $t = 0$ are

$$v_1^{(0)}(t) = \theta(-qt) \sum_{m=0}^{\infty} b_m t^m, \quad v_2^{(0)}(t) = \frac{1}{\theta(-aqt)} \sum_{m=0}^{\infty} c_m t^m.$$

We assume that $b_0 = 1$ and $c_0 = 1$. Here $u_1(t) = \sum b_m t^m$ is divergent and $u_2(t) = \sum c_m t^m$ is convergent. The q -Borel transforms of $u_1(t)$ and $u_2(t)$ are given by

$$\begin{aligned} \mathcal{B}_q^+(u_1)(\tau) &= (-a\tau; q)_\infty (-q\tau; q)_\infty, \\ \mathcal{B}_q^-(u_2)(\tau) &= \frac{1}{(-q^2\tau; q)_\infty (-aq\tau; q)_\infty}. \end{aligned} \quad (7)$$

3 q -Stokes phenomenon

We give a resummation procedure of the divergent power series $u_1(t)$ by the q -Laplace transformation and study the q -Stokes phenomenon. We set $v(t) = \theta(-qt)u(t)$ in (6). Then $u(t)$ satisfies

$$\{\sigma_q - [1 - (a + q)t] + at^2\sigma_q^{-1}\}u(t) = 0. \tag{8}$$

The series $u_1(t)$ is a unique formal power series solution of (8) around the origin with $b_0 = 1$.

The q -Borel transform of $u_1(t)$ is given by

$$\mathcal{B}_q^+(u_1)(\tau) = (-a\tau, -q\tau; q)_\infty.$$

But the q -Laplace transform of $(-a\tau, -q\tau; q)_\infty$ is divergent. We apply a q -analogue of Borel transform of order $1/2$ studied in [1, 2] in order to obtain a resummation of $u_1(t)$.

We set $p^2 = q$. We consider the p -Borel-Laplace transform of $u_1(t)$

$$f_p(t, \lambda) = L_{p;1}^{[\lambda]} \circ \mathcal{B}_p^+(u_1)(t).$$

The two choices of p give the different p -Borel-Laplace transforms. Since $p^2 = q$, $L_{p;1}^{[\lambda]}$ is considered as the p -Borel-Laplace transform of order $1/2$.

Our main result is as follows.

Theorem 4. *The p -Borel-Laplace transform $f_p(t, \lambda)$ is a meromorphic function on \mathbb{C}^* and has at most a simple pole on $t = -\lambda p^{\mathbb{Z}}$:*

$$\begin{aligned} f_p(t, \lambda) = & \frac{\theta_q(a\lambda)\theta_q(ap\lambda)}{(q/a; q)_\infty\theta_q(-ap\lambda^2)} \frac{\theta_q(-pt/\lambda^2)}{\theta_q(pt/\lambda)\theta_q(qt/\lambda)} {}_1\phi_1(0; a; q, 1/t) \\ & + \frac{\theta_q(q\lambda)\theta_q(qp\lambda)}{(a/q; q)_\infty\theta_q(-ap\lambda^2)} \frac{\theta_q(-pqt/a\lambda^2)}{\theta_q(pt/\lambda)\theta_q(qt/\lambda)} {}_1\phi_1(0; q^2/a; q, q/at). \end{aligned}$$

Proof. The divergent series $u_1(t)$ satisfies the p -difference equation

$$\{\sigma_p^2 - [1 - (a + p^2)t] + at^2\sigma_p^{-2}\}u_1(t) = 0.$$

The p -Borel transform of $\varphi(\tau) = \mathcal{B}_p^+(u_1)(\tau)$ satisfies

$$\{\sigma_p^2 + (a + p^2)\tau\sigma_p - (1 - ap\tau^2)\}\varphi(\tau) = 0. \tag{9}$$

The power series $\varphi(\tau)$ give a unique holomorphic solution around the origin of (9) with $\varphi(0) = 0$. We set $c^2 = ap$. Then $g(\tau) = (-c\tau; p)_\infty\varphi(\tau)$ satisfies

$$\{(1 + cp\tau)\sigma_p^2 + (c^2/p + p^2)\tau\sigma_p - (1 - c\tau)\}g(\tau) = 0,$$

which has a solution $g(\tau) = {}_2\phi_1(-c/p, -p^2/c; -p; p, c\tau)$. Therefore we have

$$\varphi(\tau) = \frac{1}{(-c\tau; p)_\infty} {}_2\phi_1(-c/p, -p^2/c; -p; p, c\tau).$$

We study the asymptotic behavior of $\varphi(\tau)$ around the infinity. It is evident that

$$\frac{1}{(-c\tau; p)_\infty} = \frac{(p; p)_\infty}{\theta_p(-c\tau; p)} (-p/c\tau; p)_\infty.$$

By the connection formula (3), we have

$$\begin{aligned} & {}_2\phi_1(-c/p, -p^2/c; -p; p, c\tau) \\ &= \frac{(p^2/c, -p^2/c; p)_\infty}{(-p, p^3/c^2; p)_\infty} \frac{\theta_p(c^2\tau/p)}{\theta_p(-c\tau)} {}_2\phi_1(c/p, -c/p; c^2/p^2; p, -p/c\tau) \\ &+ \frac{(c/p, -c/p; p)_\infty}{(-p, c^2/p^3; p)_\infty} \frac{\theta_p(p^2\tau)}{\theta_p(-c\tau)} {}_2\phi_1(p^2/c, -p^2/c; p^4/c^2; p, -p/c\tau). \end{aligned} \quad (10)$$

By (4), we have

$$(-p/c\tau; p)_\infty {}_2\phi_1(c/p, -c/p; c^2/p^2; p, -p/c\tau) = {}_0\phi_1(-; c^2/p; p^2, p/\tau^2).$$

$$(-p/c\tau; p)_\infty {}_2\phi_1(p^2/c, -p^2/c; p^4/c^2; p, -p/c\tau) = {}_0\phi_1(-; p^5/c^2; p^2, p^7/c^4\tau^2).$$

Therefore the behavior of $\varphi(\tau)$ at the infinity is as follows.

$$\begin{aligned} \varphi(\tau) &= \frac{(p, p^2/c, -p^2/c; p)_\infty}{(-p, p^3/c^2; p)_\infty} \frac{\theta_p(c^2\tau/p)}{\theta_p(c\tau)\theta_p(-c\tau)} {}_0\phi_1(-; c^2/p; p^2, p/\tau^2) \\ &+ \frac{(p, c/p, -c/p; p)_\infty}{(-p, c^2/p^3; p)_\infty} \frac{\theta_p(p^2\tau)}{\theta_p(c\tau)\theta_p(-c\tau)} {}_0\phi_1(-; p^5/c^2; p^2, p^7/c^4\tau^2). \end{aligned}$$

We calculate the p -Laplace transform $f_p(t, \lambda)$ of $\varphi(\tau)$ by Lemma 3:

$$\begin{aligned} f_p(t, \lambda) &= \frac{(p, p^2/c, -p^2/c; p)_\infty}{(-p, p^3/c^2; p)_\infty} \frac{\theta_p(c^2\lambda/p)\theta_{p^2}(-pt/\lambda^2)}{\theta_p(c\lambda)\theta_p(-c\lambda)\theta_p(pt/\lambda)} {}_1\phi_1(0; c^2/p; p^2, 1/t) \\ &+ \frac{(p, c/p, -c/p; p)_\infty}{(-p, c^2/p^3; p)_\infty} \frac{\theta_p(p^2\lambda)\theta_{p^2}(-p^4t/c^2\lambda^2)}{\theta_p(c\lambda)\theta_p(-c\lambda)\theta_p(pt/\lambda)} {}_1\phi_1(0; p^5/c^2; p^2, p^3/c^2t). \end{aligned}$$

By Lemma 1, we have

$$\begin{aligned} \frac{(p; p)_\infty}{(-p; p)_\infty} \frac{1}{\theta_p(c\lambda)\theta_p(-c\lambda)} &= \frac{1}{\theta_{p^2}(-c^2\lambda^2)} = \frac{1}{\theta_q(-ap\lambda^2)}, \\ \frac{(p^2/c, -p^2/c; p)_\infty}{(p^3/c^2; p)_\infty} &= \frac{(p^4/c^2; p^2)_\infty}{(p^3/c^2; p)_\infty} = \frac{1}{(p^3/c^2; p^2)_\infty} = \frac{1}{(q/a; q)_\infty}. \end{aligned}$$

And

$$\frac{\theta_p(a\lambda)\theta_q(-pt/\lambda^2)}{\theta_q(-ap\lambda^2)\theta_p(pt/\lambda)} = \frac{\theta_q(a\lambda)\theta_q(ap\lambda)\theta_q(-pt/\lambda^2)}{\theta_q(-ap\lambda^2)\theta_q(pt/\lambda)\theta_q(qt/\lambda)}.$$

Applying Lemma 1 to the second term, we obtain Theorem 4. \square

4 Connection formula of convergent series

We show a connection formula between $v_2^{(0)}(t)$ and solutions of (6) around the infinity.

Theorem 5. *The solution $v_2^{(0)}(t)$ is written by the sum of $v_1^{(\infty)}(t)$ and $v_2^{(\infty)}(t)$ on $t \in \mathbb{C}^*$:*

$$v_2^{(0)}(t) = \frac{1}{(q; q)_\infty (a/q; q)_\infty} v_1^{(\infty)}(t) + \frac{q}{a \cdot (q; q)_\infty (a/q; q)_\infty} v_2^{(\infty)}(t).$$

Remark. This relation is essentially obtained by Morita [11].

Proof. By (7), $u_2(t)$ has an integral representation

$$u_2(t) = \frac{1}{2\pi i} \int_{|\tau|=\varepsilon} \frac{1}{(-q^2\tau; q)_\infty (-aqt; q)_\infty} \theta_q(t/\tau) \frac{d\tau}{\tau},$$

by the residue calculus around the origin. Here ε is sufficiently small so that $(-q^2t; q)_\infty (-aqt; q)_\infty$ does not have zeros in $|\tau| \leq \varepsilon$.

If we take R so that the circle $|z| = R$ does not pass through the poles, we have

$$\frac{1}{2\pi i} \int_{|\tau|=R} \frac{1}{(-q^2t; q)_\infty (-aqt; q)_\infty} \theta_q(t/\tau) \frac{d\tau}{\tau} \rightarrow 0$$

when $R \rightarrow \infty$. Therefore

$$u_2(t) = - \sum_{n=0}^{\infty} \text{Res} \left\{ \frac{1}{(-q^2t; q)_\infty (-aqt; q)_\infty} \theta_q(t/\tau) \frac{d\tau}{\tau} : \tau = -q^{-n-2} \right\} \\ - \sum_{n=0}^{\infty} \text{Res} \left\{ \frac{1}{(-q^2t; q)_\infty (-aqt; q)_\infty} \theta_q(t/\tau) \frac{d\tau}{\tau} : \tau = -q^{-n-1}/a \right\}.$$

We can calculate the residues by the following lemma [18].

Lemma 6. *We assume that $b, c \in \mathbb{C}^*$, $c \notin q^{\mathbb{Z}}$ and $n = 0, 1, 2, 3, \dots$. Then we have*

$$\text{Res} \left\{ \frac{1}{(bz; q)_\infty} \frac{dz}{z} : z = q^{-n}/b \right\} = \frac{(-1)^{-n+1} q^{n(n+1)/2}}{(q; q)_\infty (q; q)_n}, \\ \theta_q(bq^n t) = q^{-n(n-1)/2} b^{-n} t^{-n} \theta_q(bt), \\ \frac{1}{(cq^{-n}; q)_\infty} = \frac{(-c)^{-n} q^{n(n+1)/2}}{(c; q)_\infty (q/c; q)_n}.$$

By the lemma above we have

$$u_2(t) = \frac{\theta_q(-q^2t)}{(q; q)_\infty (a/q; q)_\infty} {}_1\phi_1(0; a/q; q, q/at) + \frac{\theta_q(-aqt)}{(q; q)_\infty (q/a; q)_\infty} {}_1\phi_1(0; a; q, 1/t).$$

Since $u_2(t) = \theta(-aqt)v_2^{(0)}(t)$, we obtain Theorem 5. □

5 Summary

We have shown a connection formula of a second order q -difference equation whose solution is represented by ${}_1\phi_1(0; a; q, t)$:

$$qtv(tq) + [1 - (a + q)t]v(t) + atv(t/q) = 0.$$

Local solutions around $t = \infty$ are

$$v_1^{(\infty)}(t) = {}_1\phi_1(0; a; q, 1/t), \quad v_2^{(\infty)}(t) = \frac{\theta(-qt)}{\theta(-at)} {}_1\phi_1(0; q^2/a; q, q/at).$$

Local (formal) solutions around $t = 0$ are

$$v_1^{(0)}(t) = \theta(-qt) \sum_{m=0}^{\infty} b_m t^m, \quad v_2^{(0)}(t) = \frac{1}{\theta(-aqt)} \sum_{m=0}^{\infty} c_m t^m.$$

Here $\sum b_m t^m$ is divergent, $\sum c_m t^m$ is convergent. We assume that $b_0 = 1$, $c_0 = 1$. We set $\tilde{v}_1^{(0)}(t, \lambda; p) = \theta(-qt) f_p(t, \lambda)$ for $p^2 = q$. Here $f_p(t, \lambda)$ is a resummation

$$f_p(t, \lambda) = L_{p;1}^{[\lambda]} \circ \mathcal{B}_p^+ \left[\sum_{m=0}^{\infty} b_m t^m \right].$$

Theorem 7. *The connection formulae between $\tilde{v}_1^{(0)}(t, \lambda; p)$, $v_2^{(0)}(t)$ and $v_1^{(\infty)}(t)$, $v_2^{(\infty)}(t)$ are given as follows.*

$$\begin{aligned} \tilde{v}_1^{(0)}(t, \lambda; p) &= \frac{\theta_q(a\lambda)\theta_q(ap\lambda)}{(q/a; q)_\infty \theta_q(-ap\lambda^2)} \frac{\theta(-qt)\theta_q(-pt/\lambda^2)}{\theta_q(pt/\lambda)\theta_q(qt/\lambda)} v_1^{(\infty)}(t) \\ &\quad + \frac{\theta_q(q\lambda)\theta_q(qp\lambda)}{(a/q; q)_\infty \theta_q(-ap\lambda^2)} \frac{\theta(-at)\theta_q(-pqt/a\lambda^2)}{\theta_q(pt/\lambda)\theta_q(qt/\lambda)} v_2^{(\infty)}(t), \\ v_2^{(0)}(t) &= \frac{1}{(q; q)_\infty (q/a; q)_\infty} v_1^{(\infty)}(t) + \frac{q}{a \cdot (q; q)_\infty (a/q; q)_\infty} v_2^{(\infty)}(t). \end{aligned}$$

The second connection formula is already shown by Morita [11]. The case $a = -q$ is obtained in [12]. A connection formula of the Hahn-Exton q -Bessel equation is derived from the theorem above by simple calculations.

The q -Laplace transform of order $1/2$ is shown in [1, 2] is necessary to determine the q -Stokes coefficients. Our results is the first example to calculate the q -Stokes coefficients when the slope of the Newton diagram is two in q -difference equations. Our method would be useful to study the q -Stokes phenomenon of other q -difference equations with slopes higher than two.

References

- [1] Dreyfus, T., Eloy, A.; q -Borel-Laplace summation for q -difference equations with two slopes, arXiv:1501.02994 [math.CV].

- [2] Eloy, A.; Classification et géométrie des équations aux q -différences: étude globale de q -Painlevé, classification non isoformelle et Stokes à pentes arbitraires, Thèse, Université de Toulouse, 2016.
- [3] Exton, H.; A basic analogue of the Bessel-Clifford equation, *Jñānābha*, **8** (1978), 49–56.
- [4] Hahn, W.; Die mechanische Deutung einer geometrischen Differenzgleichung, *Zeitschrift für Angewandte Mathematik und Mechanik* **33** (1953), 1953 270–272.
- [5] Ismail, Mourad E. H.; The zeros of basic Bessel functions, the functions $J_{\nu+ax}(x)$, and associated orthogonal polynomials. *J. Math. Anal. Appl.* **86** (1982), 1–19.
- [6] Jackson, F. H.; On generalized functions of Legendre and Bessel, *Trans. Roy. Soc. Edinburgh*, **41** (1903), 1–28.
- [7] Jackson, F. H.; Theorems relating to a generalization of the Bessel functions, *Trans. Roy. Soc. Edinburgh*, **41** (1904), 105–118, 399–408.
- [8] Jackson, F. H.; Borel's integral and q -series, *Proc. Roy. Soc. Edinburgh*, **30** (1910), 378–385.
- [9] Gasper, G., Rahman, M.; *Basic Hypergeometric Series*, 2nd ed, Cambridge, 2004.
- [10] Hamamoto, T., Kajiwara, K., Witte, N. S.; Hypergeometric solutions to the q -Painlevé equation of type $(A_1 + A'_1)^{(1)}$, *Int. Math. Res. Not.* 2006, Art. ID 84619, 26 pp.
- [11] Morita, T.; A Connection Formula of the Hahn-Exton q -Bessel Function, *SIGMA* **7** (2011), 115, 11 pages.
- [12] Morita, T.; The Stokes phenomenon for the q -difference equation satisfied by the Ramanujan entire function, *Ramanujan J.* **34** (2014), 329–346.
- [13] Olde Daalhuis, A. B.; Asymptotic expansions for q -gamma, q -exponential, and q -Bessel functions. *J. Math. Anal. Appl.* **186** (1994), 896–913.
- [14] Ramis, J.-P., Sauloy, J., Zhang, C.; *Local analytic classification of q -difference equations*, Astérisque, 2013.
- [15] Swarttouw, R.; *The Hahn-Exton q -Bessel function*, PhD Thesis, Delft University of Technology, 1992.
- [16] Thomae J.; Ueber die Functionen welche durch Reihen von der Form dargestellt werden $1 + \frac{p}{1} \frac{p'}{q'} \frac{p''}{q''} x + \frac{p}{1} \frac{p+1}{2} \frac{p'}{q'} \frac{p'+1}{q'+1} \frac{p''}{q''} \frac{p''+1}{q''+1} x^2 + \dots$, *J. Reine Angew. Math.* **87** (1879), 26–73.

- [17] Watson, G. N., The continuation of functions defined by generalized hypergeometric series, *Trans. Camb. Phil. Soc.* **21** (1910), 281–299.
- [18] Zhang C.; Une sommation discrète pour des équations aux q -différences linéaires et à coefficients, analytiques: théorie générale et exemples, in “Differential Equations and Stokes Phenomenon”, World Sci. Publ., River Edge, NJ, 2002, 309–329.
- [19] Zhang C.; Sur les fonctions q -Bessel de Jackson, *J. Approx. Theory* **122** (2003), 208–223.
- [20] Zhang C.; Remarks on some basic hypergeometric series, in *Theory and Applications of Special Functions*, Dev. Math., Vol. 13, Springer, New York, 2005, 479–491.