

## Inequalities for the Difference $A^{-1}g(A) - B^{-1}g(B)$ when $g$ is Operator Monotone on $[0, \infty)$

By

Silvestru Sever DRAGOMIR

*College of Engineering & Science, Victoria University,  
PO Box 14428, Melbourne City, MC 8001, Australia.  
e-mail address: sever.dragomir@vu.edu.au*

(Received June 11, Revised September 14, 2020)

Abstract

In this paper we show that, if  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$  with  $g(0) = 0$ ,  $A \geq 0$  and there exist positive numbers  $d > c > 0$  such that the condition  $d1_H \geq B - A \geq c1_H > 0$  is satisfied, then

$$\begin{aligned} A^{-1}g(A) - B^{-1}g(B) &\geq \left[ \frac{g(\|A\|)}{\|A\|} - \frac{g(\|A\| + c)}{\|A\| + c} \right] 1_H \\ &\geq \left[ \frac{g(\|B\| - c)}{\|B\| - c} - \frac{g(\|B\|)}{\|B\|} \right] 1_H > 0 \end{aligned}$$

and

$$\begin{aligned} A^{-1}g(A) - B^{-1}g(B) \\ \geq c \left( \frac{g(\|A\|)}{(d + \|A\|)\|A\|} - \frac{g(d + \|A\|) - g(\|A\|)}{d(d + \|A\|)} \right) 1_H \geq 0. \end{aligned}$$

Some applications for particular functions of interest are also given.  
2010 Mathematics Subject Classification. Primary 47A63; Secondary 47A60.

## 1 Introduction

Consider a complex Hilbert space  $(H, \langle \cdot, \cdot \rangle)$ . An operator  $T$  is said to be positive (denoted by  $T \geq 0$ ) if  $\langle Tx, x \rangle \geq 0$  for all  $x \in H$  and also an operator  $T$  is said to be *strictly positive* (denoted by  $T > 0$ ) if  $T$  is positive and invertible. A real valued continuous function  $f(t)$  on  $[0, \infty)$  is said to be operator monotone if  $f(A) \geq f(B)$  holds for any  $A \geq B > 0$ .

In 1934, K. Löwner [6] had given a definitive characterization of operator monotone functions as follows, see for instance [1, p. 144-145]:

**Theorem 1** *A function  $f : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone in  $[0, \infty)$  if and only if it has the representation*

$$f(t) = f(0) + bt + \int_0^\infty \frac{ts}{t+s} dm(s) \quad (1)$$

where  $b \geq 0$  and a positive measure  $m$  on  $[0, \infty)$  such that

$$\int_0^\infty \frac{s}{1+s} dm(s) < \infty.$$

We recall the important fact proved by Löwner and Heinz that states that the power function  $f : [0, \infty) \rightarrow \mathbb{R}$ ,  $f(t) = t^\alpha$  is an operator monotone function for any  $\alpha \in [0, 1]$ , [5]. Also the function  $\ln$  is operator monotone on the open interval  $(0, \infty)$ . Let  $f(t)$  be a continuous function  $(0, \infty) \rightarrow (0, \infty)$ . It is known that  $f(t)$  is operator monotone if and only if  $g(t) = t/f(t) =: f^*(t)$  is also operator monotone, see for instance [3] or [7].

Let  $A$  and  $B$  be strictly positive operators on a Hilbert space  $H$  such that  $B - A \geq m1_H > 0$ . In 2015, [4], T. Furuta obtained the following result for any non-constant operator monotone function  $f$  on  $[0, \infty)$

$$\begin{aligned} f(B) - f(A) &\geq [f(\|A\| + m) - f(\|A\|)] 1_H \\ &\geq [f(\|B\|) - f(\|B\| - m)] 1_H > 0. \end{aligned} \quad (2)$$

If  $B > A > 0$ , then

$$\begin{aligned} f(B) - f(A) &\geq \left[ f \left( \|A\| + \frac{1}{\|(B-A)^{-1}\|} \right) - f(\|A\|) \right] 1_H \\ &\geq \left[ f(\|B\|) - f \left( \|B\| - \frac{1}{\|(B-A)^{-1}\|} \right) \right] 1_H > 0. \end{aligned} \quad (3)$$

The inequality between the first and third term in (3) was obtained earlier by H. Zuo and G. Duan in [9].

By taking  $f(t) = t^r$ ,  $r \in (0, 1]$  in (3) Furuta obtained the following refinement of the celebrated Löwner-Heinz inequality

$$\begin{aligned} B^r - A^r &\geq \left[ \left( \|A\| + \frac{1}{\|(B-A)^{-1}\|} \right)^r - \|A\|^r \right] 1_H \\ &\geq \left[ \|B\|^r - \left( \|B\| - \frac{1}{\|(B-A)^{-1}\|} \right)^r \right] 1_H > 0 \end{aligned} \quad (4)$$

provided  $B > A > 0$ .

With the same assumptions for  $A$  and  $B$ , we have the logarithmic inequality [4]

$$\begin{aligned} \ln B - \ln A &\geq \left[ \ln \left( \|A\| + \frac{1}{\|(B-A)^{-1}\|} \right) - \ln(\|A\|) \right] 1_H \\ &\geq \left[ \ln(\|B\|) - \ln \left( \|B\| - \frac{1}{\|(B-A)^{-1}\|} \right) \right] 1_H > 0. \end{aligned} \quad (5)$$

Notice that the inequalities between the first and third terms in (4) and (5) were obtained earlier by M. S. Moslehian and H. Najafi in [8].

Motivated by the above results, in this paper we show that, if  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$  with  $g(0) = 0$ ,  $A \geq 0$  and there exist positive numbers  $d > c > 0$  such that the condition  $d1_H \geq B - A \geq c1_H > 0$  is satisfied, then

$$\begin{aligned} A^{-1}g(A) - B^{-1}g(B) &\geq \left[ \frac{g(\|A\|)}{\|A\|} - \frac{g(\|A\| + c)}{\|A\| + c} \right] 1_H \\ &\geq \left[ \frac{g(\|B\| - c)}{\|B\| - c} - \frac{g(\|B\|)}{\|B\|} \right] 1_H > 0 \end{aligned}$$

and

$$\begin{aligned} &A^{-1}g(A) - B^{-1}g(B) \\ &\geq c \left( \frac{g(\|A\|)}{(d + \|A\|)\|A\|} - \frac{g(d + \|A\|) - g(\|A\|)}{d(d + \|A\|)} \right) 1_H \geq 0. \end{aligned}$$

Some applications for particular functions of interest are also given.

## 2 Main Results

We start with the following lemma that is of interest in itself.

**Lemma 2** *Assume that  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . Then the function  $f : (0, \infty) \rightarrow \mathbb{R}$ ,*

$$f(t) := \frac{g(0) - g(t)}{t} \quad (6)$$

*is operator monotone on  $(0, \infty)$ . If  $g(0) = 0$ , then  $f(t) = -g(t)t^{-1}$  is operator monotone on  $(0, \infty)$ .*

**Proof.** Since  $g$  is operator monotone on  $[0, \infty)$ , then there exists  $b \geq 0$  and  $w$  is a positive measure satisfying

$$\int_0^\infty \frac{\lambda}{1 + \lambda} dw(\lambda) < \infty$$

such that [1, p. 144-145]

$$g(t) = g(0) + bt + \int_0^\infty \frac{\lambda t}{t + \lambda} dw(\lambda). \quad (7)$$

We have for  $t > 0$  that

$$h(t) := \frac{g(t) - g(0)}{t} - b = \int_0^\infty \frac{\lambda}{t + \lambda} dw(\lambda)$$

Therefore for all  $A, B > 0$

$$h(B) - h(A) = \int_0^\infty \lambda \left[ (B + \lambda 1_H)^{-1} - (A + \lambda 1_H)^{-1} \right] dw(\lambda). \quad (8)$$

Let  $T, S > 0$ . The function  $g(t) = -t^{-1}$  is operator monotonic on  $(0, \infty)$ , operator Gâteaux differentiable and the Gâteaux derivative is given by

$$\nabla g_T(S) := \lim_{t \rightarrow 0} \left[ \frac{g(T + tS) - g(T)}{t} \right] = T^{-1} S T^{-1} \quad (9)$$

for  $T, S > 0$ .

Consider the continuous function  $g$  defined on an interval  $I$  for which the corresponding operator function is Gâteaux differentiable and for  $C, D$  selfadjoint operators with spectra in  $I$  we consider the auxiliary function defined on  $[0, 1]$  by

$$g_{C,D}(t) := g((1-t)C + tD), \quad t \in [0, 1].$$

Then we have, by the properties of the Bochner integral, that

$$g(D) - g(C) = \int_0^1 \frac{d}{dt} (g_{C,D}(t)) dt = \int_0^1 \nabla g_{(1-t)C+tD}(D - C) dt. \quad (10)$$

If we write this equality for the function  $g(t) = -t^{-1}$  and  $C, D > 0$ , then we get the representation

$$C^{-1} - D^{-1} = \int_0^1 ((1-t)C + tD)^{-1} (D - C) ((1-t)C + tD)^{-1} dt. \quad (11)$$

Now, if we replace in (11)  $C = B + \lambda 1_H$  and  $D = A + \lambda 1_H$  for  $\lambda > 0$ , then we get

$$\begin{aligned} & (B + \lambda 1_H)^{-1} - (A + \lambda 1_H)^{-1} \\ &= \int_0^1 ((1-t)B + tA + \lambda 1_H)^{-1} (A - B) ((1-t)B + tA + \lambda 1_H)^{-1} dt. \end{aligned} \quad (12)$$

Therefore, by (8),

$$\begin{aligned}
 h(B) - h(A) &= \int_0^\infty \lambda \left( \int_0^1 ((1-t)B + tA + \lambda 1_H)^{-1} (A - B) \right. \\
 &\quad \left. \times ((1-t)B + tA + \lambda 1_H)^{-1} dt \right) dw(\lambda) \\
 &= - \int_0^\infty \lambda \left( \int_0^1 ((1-t)B + tA + \lambda 1_H)^{-1} (B - A) \right. \\
 &\quad \left. \times ((1-t)B + tA + \lambda 1_H)^{-1} dt \right) dw(\lambda).
 \end{aligned} \tag{13}$$

If  $B \geq A > 0$ , then

$$((1-t)B + tA + \lambda 1_H)^{-1} (B - A) ((1-t)B + tA + \lambda 1_H)^{-1} \geq 0$$

for all  $t, \lambda > 0$ , which implies that

$$\begin{aligned}
 &\int_0^\infty \lambda \left( \int_0^1 ((1-t)B + tA + \lambda 1_H)^{-1} (B - A) \right. \\
 &\quad \left. \times ((1-t)B + tA + \lambda 1_H)^{-1} dt \right) dw(\lambda) \geq 0,
 \end{aligned}$$

namely

$$\begin{aligned}
 f(B) - f(A) &= h(A) - h(B) \\
 &= \int_0^\infty \lambda \left( \int_0^1 ((1-t)B + tA + \lambda 1_H)^{-1} (B - A) \right. \\
 &\quad \left. \times ((1-t)B + tA + \lambda 1_H)^{-1} dt \right) dw(\lambda) \\
 &\geq 0.
 \end{aligned}$$

Therefore, the function  $f$  is operator monotone on  $(0, \infty)$ . ■

**Theorem 3** Assume that  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . If  $A > 0$  and there exists  $c > 0$  such that  $B - A \geq c1_H > 0$ , then

$$\begin{aligned}
 &A^{-1}g(A) - B^{-1}g(B) - g(0)(A^{-1} - B^{-1}) \\
 &\geq \left[ \frac{g(\|A\|)}{\|A\|} - \frac{g(\|A\| + c)}{\|A\| + c} - g(0) \frac{c}{(\|A\| + c)\|A\|} \right] 1_H \\
 &\geq \left[ \frac{g(\|B\| - c)}{\|B\| - c} - \frac{g(\|B\|)}{\|B\|} - g(0) \frac{c}{(\|B\| - c)\|B\|} \right] 1_H > 0.
 \end{aligned} \tag{14}$$

If  $g(0) = 0$ , then

$$\begin{aligned}
 A^{-1}g(A) - B^{-1}g(B) &\geq \left[ \frac{g(\|A\|)}{\|A\|} - \frac{g(\|A\| + c)}{\|A\| + c} \right] 1_H \\
 &\geq \left[ \frac{g(\|B\| - c)}{\|B\| - c} - \frac{g(\|B\|)}{\|B\|} \right] 1_H > 0.
 \end{aligned} \tag{15}$$

**Proof.** If we write the inequality (2) for  $f(t) = \frac{g(0)-g(t)}{t}$ ,  $t > 0$ , which, by Lemma 2, is operator monotone, then we have

$$\begin{aligned} & B^{-1} [g(0) - g(B)] - A^{-1} [g(0) - g(A)] \\ & \geq \left[ \frac{g(0) - g(\|A\| + c)}{\|A\| + c} - \frac{g(0) - g(\|A\|)}{\|A\|} \right] \mathbf{1}_H \\ & \geq \left[ \frac{g(0) - g(\|B\|)}{\|B\|} - \frac{g(0) - g(\|B\| - c)}{\|B\| - c} \right] \mathbf{1}_H > 0. \end{aligned} \quad (16)$$

Observe that

$$\begin{aligned} & B^{-1} [g(0) - g(B)] - A^{-1} [g(0) - g(A)] \\ & = A^{-1}g(A) - B^{-1}g(B) - g(0) (A^{-1} - B^{-1}), \\ & \frac{g(0) - g(\|A\| + c)}{\|A\| + c} - \frac{g(0) - g(\|A\|)}{\|A\|} \\ & = \frac{g(\|A\|)}{\|A\|} - \frac{g(\|A\| + c)}{\|A\| + c} - g(0) \frac{c}{(\|A\| + c)\|A\|} \end{aligned}$$

and

$$\begin{aligned} & \frac{g(0) - g(\|B\|)}{\|B\|} - \frac{g(0) - g(\|B\| - c)}{\|B\| - c} \\ & = \frac{g(\|B\| - c)}{\|B\| - c} - \frac{g(\|B\|)}{\|B\|} - g(0) \frac{c}{(\|B\| - c)\|B\|} \end{aligned}$$

and by (16) we get (14). ■

Its is well known that, if  $P \geq 0$ , then

$$|\langle Px, y \rangle|^2 \leq \langle Px, x \rangle \langle Py, y \rangle$$

for all  $x, y \in H$ .

Therefore, if  $T > 0$ , then

$$\begin{aligned} 0 & \leq \langle x, x \rangle^2 = \langle T^{-1}Tx, x \rangle^2 = \langle Tx, T^{-1}x \rangle^2 \\ & \leq \langle Tx, x \rangle \langle TT^{-1}x, T^{-1}x \rangle = \langle Tx, x \rangle \langle x, T^{-1}x \rangle \end{aligned}$$

for all  $x \in H$ .

If  $x \in H$ ,  $\|x\| = 1$ , then

$$1 \leq \langle Tx, x \rangle \langle x, T^{-1}x \rangle \leq \langle Tx, x \rangle \sup_{\|x\|=1} \langle x, T^{-1}x \rangle = \langle Tx, x \rangle \|T^{-1}\|,$$

which implies the following operator inequality

$$\|T^{-1}\|^{-1} \mathbf{1}_H \leq T. \quad (17)$$

**Corollary 4** Assume that  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned}
 & A^{-1}g(A) - B^{-1}g(B) - g(0)(A^{-1} - B^{-1}) \tag{18} \\
 & \geq \left[ \frac{g(\|A\|)}{\|A\|} - \frac{g\left(\|A\| + \|(B-A)^{-1}\|^{-1}\right)}{\|A\| + \|(B-A)^{-1}\|^{-1}} \right] 1_H \\
 & \quad - g(0) \frac{\|(B-A)^{-1}\|^{-1}}{\left(\|A\| + \|(B-A)^{-1}\|^{-1}\right) \|A\|} 1_H \\
 & \geq \left[ \frac{g\left(\|B\| - \|(B-A)^{-1}\|^{-1}\right)}{\|B\| - \|(B-A)^{-1}\|^{-1}} - \frac{g(\|B\|)}{\|B\|} \right] 1_H \\
 & \quad - g(0) \frac{\|(B-A)^{-1}\|^{-1}}{\left(\|B\| - \|(B-A)^{-1}\|^{-1}\right) \|B\|} 1_H \\
 & > 0.
 \end{aligned}$$

If  $g(0) = 0$ , then

$$\begin{aligned}
 & A^{-1}g(A) - B^{-1}g(B) \tag{19} \\
 & \geq \left[ \frac{g(\|A\|)}{\|A\|} - \frac{g\left(\|A\| + \|(B-A)^{-1}\|^{-1}\right)}{\|A\| + \|(B-A)^{-1}\|^{-1}} \right] 1_H \\
 & \geq \left[ \frac{g\left(\|B\| - \|(B-A)^{-1}\|^{-1}\right)}{\|B\| - \|(B-A)^{-1}\|^{-1}} - \frac{g(\|B\|)}{\|B\|} \right] 1_H > 0.
 \end{aligned}$$

We have the following lower bound as well:

**Theorem 5** Assume that  $f : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . Let  $A \geq 0$  and assume that there exist positive numbers  $d > c > 0$  such that

$$d1_H \geq B - A \geq c1_H > 0. \tag{20}$$

Then

$$f(B) - f(A) \geq c \frac{f(d + \|A\|) - f(\|A\|)}{d} 1_H \geq 0. \tag{21}$$

**Proof.** Since the function  $f : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone in  $[0, \infty)$ , then  $f$  can be written as in the equation (1) and for  $A, B \geq 0$  we have the representation

$$\begin{aligned} f(B) - f(A) & \qquad \qquad \qquad (22) \\ & = b(B - A) + \int_0^\infty s \left[ B(B + s1_H)^{-1} - A(A + s1_H)^{-1} \right] dm(s). \end{aligned}$$

Observe that for  $s > 0$

$$\begin{aligned} & B(B + s1_H)^{-1} - A(A + s1_H)^{-1} \\ & = (B + s1_H - s1_H)(B + s1_H)^{-1} - (A + s1_H - s1_H)(A + s1_H)^{-1} \\ & = (B + s1_H)(B + s1_H)^{-1} - s1_H(B + s1_H)^{-1} \\ & \quad - (A + s1_H)(A + s1_H)^{-1} + s1_H(A + s1_H)^{-1} \\ & = 1_H - s1_H(B + s1_H)^{-1} - 1_H + s1_H(A + s1_H)^{-1} \\ & = s \left[ (A + s1_H)^{-1} - (B + s1_H)^{-1} \right]. \end{aligned}$$

Therefore, (22) becomes, see also [4]

$$f(B) - f(A) = b(B - A) + \int_0^\infty s^2 \left[ (A + s1_H)^{-1} - (B + s1_H)^{-1} \right] dm(s). \quad (23)$$

Now, if we replace in (11)  $C = A + s1_H$  and  $D = B + s1_H$  for  $s > 0$ , then we get

$$\begin{aligned} & (A + s1_H)^{-1} - (B + s1_H)^{-1} & (24) \\ & = \int_0^1 ((1-t)A + tB + s1_H)^{-1} (B - A) ((1-t)A + tB + s1_H)^{-1} dt. \end{aligned}$$

By the representation (23), we derive the following identity of interest

$$\begin{aligned} f(B) - f(A) & = b(B - A) & (25) \\ & + \int_0^\infty s^2 \left[ \int_0^1 ((1-t)A + tB + s1_H)^{-1} \right. \\ & \quad \left. \times (B - A) ((1-t)A + tB + s1_H)^{-1} dt \right] dm(s) \end{aligned}$$

for  $A, B > 0$ .

From the representation (25) we get for  $B = x1_H, A = 0$  that

$$f(x) - f(0) - bx = \int_0^\infty s^2 \left( \int_0^1 (tx + s1_H)^{-1} x (tx + s1_H)^{-1} dt \right) dm(s),$$

which gives for  $x > 0$  that

$$\frac{f(x) - f(0)}{x} - b = \int_0^\infty s^2 \left( \int_0^1 (tx + s)^{-2} dt \right) dm(s). \quad (26)$$



Since  $0 < c1_H \leq B - A$ , hence

$$\begin{aligned} & c((1-t)A + tB + s1_H)^{-2} \\ & \leq ((1-t)A + tB + s1_H)^{-1} (B - A) ((1-t)A + tB + s1_H)^{-1} \end{aligned}$$

for  $t \in [0, 1]$  and  $s > 0$  and by (25) we get

$$\begin{aligned} & c \int_0^\infty s^2 \left( \int_0^1 ((1-t)A + tB + s1_H)^{-2} dt \right) dm(s) \\ & \leq f(B) - f(A) - b(B - A). \end{aligned} \quad (27)$$

Observe that for  $t \in [0, 1]$  and  $s > 0$ , we have

$$\begin{aligned} (1-t)A + tB + s1_H &= A + t(B - A) + s1_H \leq A + td1_H + s1_H \\ &= (1-t)A + t(d1_H + A) + s1_H. \end{aligned}$$

Since  $A \leq \|A\| 1_H$  then

$$(1-t)A + t(d1_H + A) + s1_H \leq ((1-t)\|A\| + t(d + \|A\|) + s)1_H,$$

which implies that

$$(1-t)A + tB + s1_H \leq ((1-t)\|A\| + t(d + \|A\|) + s)1_H$$

for  $t \in [0, 1]$  and  $s > 0$ .

This implies that

$$((1-t)A + tB + s1_H)^{-1} \geq ((1-t)\|A\| + t(d + \|A\|) + s)^{-1} 1_H$$

and

$$((1-t)A + tB + s1_H)^{-2} \geq ((1-t)\|A\| + t(d + \|A\|) + s)^{-2} 1_H$$

for  $t \in [0, 1]$  and  $s > 0$ .

Therefore

$$\begin{aligned} & \int_0^\infty s^2 \left( \int_0^1 ((1-t)A + tB + s1_H)^{-2} dt \right) dm(s) \\ & \geq \int_0^\infty s^2 \left( \int_0^1 ((1-t)\|A\| + t(d + \|A\|) + s)^{-2} dt \right) dm(s) 1_H (\geq 0) \\ & = \frac{1}{d} \int_0^\infty s^2 \left( \int_0^1 ((1-t)\|A\| + t(d + \|A\|) + s)^{-1} (d + \|A\| - \|A\|) \right. \\ & \quad \times ((1-t)\|A\| + t(d + \|A\|) + s)^{-1} dt \left. \right) dm(s) 1_H \\ & = \frac{1}{d} [(f(d + \|A\|) - f(\|A\|) - bd)] 1_H \text{ (by identity (26))} \\ & = \left( \frac{f(d + \|A\|) - f(\|A\|)}{d} - b \right) 1_H \geq 0. \end{aligned}$$

By (27) we get

$$\begin{aligned}
 & f(B) - f(A) - b(B - A) \\
 & \geq c \int_0^\infty s^2 \left( \int_0^1 ((1-t)A + tB + s1_H)^{-2} dt \right) dm(s) \\
 & \geq c \left( \frac{f(d + \|A\|) - f(\|A\|)}{d} - b \right) 1_H \geq 0.
 \end{aligned} \tag{28}$$

From (28) we derive

$$\begin{aligned}
 f(B) - f(A) & \geq b(B - A) + c \left( \frac{f(d + \|A\|) - f(\|A\|)}{d} - b \right) 1_H \\
 & = b[(B - A) - c] + c \frac{f(d + \|A\|) - f(\|A\|)}{d} 1_H \\
 & \geq c \frac{f(d + \|A\|) - f(\|A\|)}{d} 1_H \geq 0
 \end{aligned}$$

since  $b[(B - A) - c] \geq 0$  and the inequality (21) is obtained. ■

**Corollary 6** Assume that  $f : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . If  $A \geq 0$  and  $B - A > 0$ , then

$$\begin{aligned}
 f(B) - f(A) & \geq \frac{f(\|B - A\| + \|A\|) - f(\|A\|)}{\|(B - A)^{-1}\| \|B - A\|} 1_H \\
 & \geq \frac{f(\|B\|) - f(\|A\|)}{\|(B - A)^{-1}\| \|B - A\|} 1_H \geq 0.
 \end{aligned} \tag{29}$$

The first inequality follows by (21) for  $d = \|B - A\|$  and  $c = \|(B - A)^{-1}\|^{-1}$ . The second and third inequalities are obvious.

**Theorem 7** Assume that  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . If  $A > 0$  and there exist positive numbers  $d > c > 0$  such that the condition (20) is satisfied, then

$$\begin{aligned}
 & g(0) (B^{-1} - A^{-1}) + A^{-1}g(A) - B^{-1}g(B) \\
 & \geq c \left( \frac{g(\|A\|) - g(0)}{(d + \|A\|) \|A\|} - \frac{g(d + \|A\|) - g(\|A\|)}{d(d + \|A\|)} \right) 1_H \geq 0.
 \end{aligned} \tag{30}$$

If  $g(0) = 0$ , then

$$\begin{aligned}
 & A^{-1}g(A) - B^{-1}g(B) \\
 & \geq c \left( \frac{g(\|A\|)}{(d + \|A\|) \|A\|} - \frac{g(d + \|A\|) - g(\|A\|)}{d(d + \|A\|)} \right) 1_H \geq 0.
 \end{aligned} \tag{31}$$

**Proof.** Since  $g$  is operator monotone, then by Lemma 2 the function  $f(t) := \frac{g(0)-g(t)}{t}$  is operator monotone on  $(0, \infty)$  and by (21) we obtain

$$\frac{g(0) - g(B)}{B} - \frac{g(0) - g(A)}{A} \geq c \frac{\frac{g(0)-g(d+\|A\|)}{d+\|A\|} - \frac{g(0)-g(\|A\|)}{\|A\|}}{d} 1_H \geq 0. \quad (32)$$

Observe that

$$\begin{aligned} & \frac{g(0) - g(B)}{B} - \frac{g(0) - g(A)}{A} \\ &= g(0) (B^{-1} - A^{-1}) + A^{-1}g(A) - B^{-1}g(B) \end{aligned}$$

and

$$\begin{aligned} & \frac{g(0) - g(d + \|A\|)}{d + \|A\|} - \frac{g(0) - g(\|A\|)}{\|A\|} \\ &= \frac{[g(0) - g(d + \|A\|)] \|A\| - [g(0) - g(\|A\|)] (d + \|A\|)}{(d + \|A\|) \|A\|} \\ &= \frac{g(0) \|A\| - g(d + \|A\|) \|A\| - g(0) d + g(\|A\|) d - g(0) \|A\| + g(\|A\|) \|A\|}{(d + \|A\|) \|A\|} \\ &= \frac{g(\|A\|) d - g(0) d + g(\|A\|) \|A\| - g(d + \|A\|) \|A\|}{(d + \|A\|) \|A\|} \\ &= d \frac{g(\|A\|) - g(0)}{(d + \|A\|) \|A\|} - \frac{g(d + \|A\|) - g(\|A\|)}{(d + \|A\|)}, \end{aligned}$$

which gives

$$\frac{\frac{g(0)-g(d+\|A\|)}{d+\|A\|} - \frac{g(0)-g(\|A\|)}{\|A\|}}{d} = \frac{g(\|A\|) - g(0)}{(d + \|A\|) \|A\|} - \frac{g(d + \|A\|) - g(\|A\|)}{d(d + \|A\|)}.$$

Then by (32) we get (30). ■

**Corollary 8** Assume that  $g : [0, \infty) \rightarrow \mathbb{R}$  is operator monotone on  $[0, \infty)$ . If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned} & g(0) (B^{-1} - A^{-1}) + A^{-1}g(A) - B^{-1}g(B) \quad (33) \\ & \geq \left\| (B - A)^{-1} \right\|^{-1} \\ & \times \left( \frac{g(\|A\|) - g(0)}{(\|B - A\| + \|A\|) \|A\|} - \frac{g(\|B - A\| + \|A\|) - g(\|A\|)}{\|B - A\| (\|B - A\| + \|A\|)} \right) 1_H \\ & \geq 0. \end{aligned}$$

If  $g(0) = 0$ , then

$$\begin{aligned}
& A^{-1}g(A) - B^{-1}g(B) \\
& \geq \left\| (B - A)^{-1} \right\|^{-1} \\
& \times \left( \frac{g(\|A\|)}{(\|B - A\| + \|A\|)\|A\|} - \frac{g(\|B - A\| + \|A\|) - g(\|A\|)}{\|B - A\|(\|B - A\| + \|A\|)} \right) 1_H \\
& \geq 0.
\end{aligned} \tag{34}$$

### 3 Some Examples

Consider the function  $g(t) = t^r$ ,  $r \in (0, 1]$ . This function is operator monotone and by (15) we have

$$\begin{aligned}
A^{r-1} - B^{r-1} & \geq \left[ \|A\|^{r-1} - (\|A\| + c)^{r-1} \right] 1_H \\
& \geq \left[ (\|B\| - c)^{r-1} - \|B\|^{r-1} \right] 1_H > 0
\end{aligned} \tag{35}$$

provided that  $A > 0$  and  $B - A \geq c1_H > 0$ .

If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned}
A^{r-1} - B^{r-1} & \geq \left[ \|A\|^{r-1} - \left( \|A\| + \left\| (B - A)^{-1} \right\|^{-1} \right)^{r-1} \right] 1_H \\
& \geq \left[ \left( \|B\| - \left\| (B - A)^{-1} \right\|^{-1} \right)^{r-1} - \|B\|^{r-1} \right] 1_H > 0.
\end{aligned} \tag{36}$$

From (21) we obtain

$$B^r - A^r \geq c \frac{(d + \|A\|)^r - \|A\|^r}{d} 1_H \geq 0 \tag{37}$$

provided that there exist positive numbers  $d > c > 0$  such that condition (20) is satisfied.

If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned}
B^r - A^r & \geq \left\| (B - A)^{-1} \right\|^{-1} \frac{(\|B - A\| + \|A\|)^r - \|A\|^r}{\|B - A\|} 1_H \\
& \geq \left\| (B - A)^{-1} \right\|^{-1} \frac{\|B\|^r - \|A\|^r}{\|B - A\|} 1_H \geq 0.
\end{aligned} \tag{38}$$

From (30) we have

$$\begin{aligned}
& A^{r-1} - B^{r-1} \\
& \geq c \left( \frac{\|A\|^{r-1}}{d + \|A\|} - \frac{(d + \|A\|)^r - \|A\|^r}{d(d + \|A\|)} \right) 1_H \geq 0,
\end{aligned} \tag{39}$$

provided that there exist positive numbers  $d > c > 0$  such that condition (20) is satisfied.

If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned} A^{r-1} - B^{r-1} &\geq \left\| (B - A)^{-1} \right\|^{-1} \\ &\times \left( \frac{\|A\|^{r-1}}{\|B - A\| + \|A\|} - \frac{(\|B - A\| + \|A\|)^r - \|A\|^r}{\|B - A\| (\|B - A\| + \|A\|)} \right) 1_H \\ &\geq 0. \end{aligned} \quad (40)$$

Consider the function  $g(t) = \ln(t + 1)$ , which is operator monotone on  $[0, \infty)$  and  $g(0) = 0$ . By Lemma 2 we get that the function  $f(t) = -t^{-1} \ln(t + 1)$  is operator monotone on  $(0, \infty)$ .

From (15) we get

$$\begin{aligned} &A^{-1} \ln(A + 1_H) - B^{-1} \ln(B + 1_H) \\ &\geq \left( \frac{\ln(\|A\| + 1)}{\|A\|} - \frac{\ln(\|A\| + 1 + c)}{\|A\| + c} \right) 1_H \\ &\geq \left( \frac{\ln(\|B\| + 1 - c)}{\|B\| - c} - \frac{\ln(\|B\| + 1)}{\|B\|} \right) 1_H > 0 \end{aligned} \quad (41)$$

provided that  $A > 0$  and  $B - A \geq c1_H > 0$ .

If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned} &A^{-1} \ln(A + 1_H) - B^{-1} \ln(B + 1_H) \\ &\geq \left( \frac{\ln(\|A\| + 1)}{\|A\|} - \frac{\ln\left(\|A\| + 1 + \left\| (B - A)^{-1} \right\|^{-1}\right)}{\|A\| + \left\| (B - A)^{-1} \right\|^{-1}} \right) 1_H \\ &\geq \left( \frac{\ln\left(\|B\| + 1 - \left\| (B - A)^{-1} \right\|^{-1}\right)}{\|B\| - \left\| (B - A)^{-1} \right\|^{-1}} - \frac{\ln(\|B\| + 1)}{\|B\|} \right) 1_H > 0. \end{aligned} \quad (42)$$

From (21) we derive

$$\ln(B + 1_H) - \ln(A + 1_H) \geq c \frac{\ln(d + \|A\| + 1) - \ln(\|A\| + 1)}{d} 1_H \geq 0 \quad (43)$$

provided that there exist positive numbers  $d > c > 0$  such that the condition (20) is satisfied.

If  $A > 0$  and  $B - A > 0$ , then

$$\begin{aligned} &\ln(B + 1_H) - \ln(A + 1_H) \\ &\geq \left\| (B - A)^{-1} \right\|^{-1} \frac{\ln(\|B - A\| + \|A\| + 1) - \ln(\|A\| + 1)}{\|B - A\|} 1_H \\ &\geq \left\| (B - A)^{-1} \right\|^{-1} \frac{\ln(\|B\| + 1) - \ln(\|A\| + 1)}{\|B - A\|} 1_H \geq 0. \end{aligned} \quad (44)$$

From (30) we have

$$\begin{aligned} & A^{-1} \ln(A+1) - B^{-1} \ln(B+1) \\ & \geq c \left( \frac{\ln(\|A\|+1)}{(d+\|A\|)\|A\|} - \frac{\ln(d+\|A\|+1) - \ln(\|A\|+1)}{d(d+\|A\|)} \right) 1_H \geq 0 \end{aligned} \quad (45)$$

provided that there exist positive numbers  $d > c > 0$  such that the condition (20) is satisfied.

Finally, from (34) we derive

$$\begin{aligned} & A^{-1} \ln(A+1_H) - B^{-1} \ln(B+1_H) \\ & \geq \left\| (B-A)^{-1} \right\|^{-1} \\ & \times \left( \frac{\ln(\|A\|+1)}{(\|B-A\|+\|A\|)\|A\|} - \frac{\ln(\|B-A\|+\|A\|+1) - \ln(\|A\|+1)}{\|B-A\|(\|B-A\|+\|A\|)} \right) 1_H \\ & \geq 0, \end{aligned} \quad (46)$$

provided  $A > 0$  and  $B - A > 0$ .

## References

- [1] R. Bhatia, Matrix Analysis. Graduate Texts in Mathematics, 169. Springer-Verlag, New York, 1997. xii+347 pp. ISBN: 0-387-94846-5.
- [2] J. I. Fujii, Y. Seo, On Parametrized Operator Means Dominated by Power Ones, *Sci. Math.* **1** (1998) 301–306.
- [3] T. Furuta, Concrete Examples of Operator Monotone Functions Obtained by an Elementary Method Without Appealing to Löwner Integral Representation, *Linear Algebra and its Applications* **429** (2008) 972–980.
- [4] T. Furuta, Precise Lower Bound of  $f(A) - f(B)$  for  $A > B > 0$  and Non-constant Operator Monotone Function  $f$  on  $[0, \infty)$ . *J. Math. Inequal.* **9** (2015), no. 1, 47–52.
- [5] E. Heinz, Beiträge zur Störungstheorie der Spektralzerlegung, *Math. Ann.* **123** (1951) 415–438.
- [6] K. Löwner, Über Monotone MatrixFunktionen, *Math. Z.* **38** (1934) 177–216.
- [7] F. Kubo, T. Ando, Means of Positive Linear Operators, *Math. Ann.* **246** (1980) 205–224.
- [8] M. S. Moslehian, H. Najafi, An Extension of the Löwner-Heinz Inequality, *Linear Algebra Appl.*, **437** (2012), 2359–2365.
- [9] H. Zuo, G. Duan, Some Inequalities of Operator Monotone Functions. *J. Math. Inequal.* **8** (2014), no. 4, 777–781.