

Asymptotic Properties of Solutions for Lanchester Type Models with Time Dependent Coefficients

By

Kosuke ONO

*Department of Mathematical Sciences,
Tokushima University, Tokushima 770-8502, JAPAN*

e-mail : k.ono@tokushima-u.ac.jp

(Received September 30, 2023)

Abstract

We consider an ordinary differential system which is a so-called Lanchester's linear law model with time dependent coefficients. We study on asymptotic forms of solutions that decay to a point on the x -axis and y -axis.

2010 Mathematics Subject Classification. 34A34, 34D05

1 Introduction

In this paper, we consider the ordinary differential system of the form :

$$\begin{cases} x'(t) = -a(t)x(t)y(t) \\ y'(t) = -b(t)x(t)y(t) \end{cases} \quad (1.1)$$

where $a(t)$ and $b(t)$ are positive continuous functions on $[0, \infty)$, and satisfy

$$A(t) = \int_0^t a(s) ds \rightarrow \infty \quad \text{and} \quad B(t) = \int_0^t b(s) ds \rightarrow \infty \quad (1.2)$$

as $t \rightarrow \infty$.

The initial value problem (1.1) with positive initial data

$$x(0) = x_0 > 0 \quad \text{and} \quad y(0) = y_0 > 0 \quad (1.3)$$

has non-negative solutions.

System (1.1) is known as one of Lanchester type models, which describes many phenomena appearing in economics, logistics, biology, and so on.

In [5], F.W.Lanchester first proposed system (1.1) to describe combat situations (see Taylor [9] for a review). System (1.1) is said a model of guerrilla engagements (see [2], [3], [10] and the references cited therein).

There are some mathematically treated research works for Lanchester type models (see [4] and [11] for Lanchester linear-law models, and see [6] and [8] for Lanchester square-law models, and also see [1] and [7] for Lanchester models with mixed forces).

First, we consider the special case where $a(t) = \alpha$ and $b(t) = \beta$ for some positive constants $\alpha > 0$ and $\beta > 0$, that is,

$$\begin{cases} x'(t) = -\alpha x(t)y(t) \\ y'(t) = -\beta x(t)y(t) \end{cases} \quad (1.4)$$

with initial data (1.3), then we have that $A(t) = \alpha t$ and $B(t) = \beta t$ in (1.2). Using the exchange ratio $E = \alpha/\beta$ of (1.4), we can easily see that $(x(t) - Ey(t))' = 0$, and hence, $x(t) - Ey(t)$ is a constant value which is denoted by symbol M , that is,

$$x(t) - Ey(t) = x_0 - Ey_0 = M. \quad (1.5)$$

Thus, we have from (1.4) and (1.5) that

$$x'(t) = -\alpha x(t)y(t) = -\beta x(t)(x(t) - M),$$

and moreover, by fundamental calculation we obtain the following representation formula of solution $(x(t), y(t))$ of (1.4) :

(1) When $M = 0$ (i.e. $x_0 = Ey_0$),

$$x(t) = (x_0^{-1} + \beta t)^{-1} \quad \text{and} \quad y(t) = (y_0^{-1} + \alpha t)^{-1} \quad (1.6)$$

for $t \geq 0$.

(2) When $M \neq 0$ (i.e. $x_0 \neq Ey_0$),

$$x(t) = \frac{M}{1 - (1 - M/x_0)e^{-M\beta t}} = \frac{(x_0/(x_0 + EN))ENe^{-EN\beta t}}{(1 - (x_0/(x_0 + EN))e^{-EN\beta t})} \quad (1.7)$$

and

$$y(t) = \frac{(y_0/(y_0 + M/E))(M/E)e^{-(M/E)\alpha t}}{1 - (y_0/(y_0 + M/E))e^{-(M/E)\alpha t}} = \frac{N}{1 - (1 - (N/y_0))e^{-N\alpha t}} \quad (1.8)$$

where $N = -M/E$ and hence

$$x(t) - M = \frac{(1 - M/x_0)e^{-M\beta t}}{1 - (1 - M/x_0)e^{-M\beta t}} \quad (1.9)$$

and

$$y(t) - N = \frac{(1 - N/y_0)e^{-N\alpha t}}{1 - (1 - N/y_0)e^{-N\alpha t}} \tag{1.10}$$

for $t \geq 0$.

In what follows, " $f(t) \sim g(t)$ as $t \rightarrow \infty$ " means that $\lim_{t \rightarrow \infty} f(t)/g(t) = 1$ for positive functions $f(t)$ and $g(t)$ defined near $+\infty$. Similarly, for vector-valued functions " $(f_1(t), f_2(t)) \sim (g_1(t), g_2(t))$ as $t \rightarrow \infty$ " means that $f_i(t) \sim g_i(t)$ as $t \rightarrow \infty$, $i = 1, 2$.

Immediately, we can obtain from (1.6)–(1.10) the following decay properties of solution $(x(t), y(t))$ of (1.4) :

(i) When $M = 0$, $(x(t), y(t)) \rightarrow (0, 0)$ as $t \rightarrow \infty$ and

$$(x(t), y(t)) \sim ((\beta t)^{-1}, (\alpha t)^{-1}) \text{ as } t \rightarrow \infty. \tag{1.11}$$

(ii) When $M > 0$, $(x(t), y(t)) \rightarrow (M, 0)$ as $t \rightarrow \infty$ and

$$(\log(x(t) - M), \log y(t)) \sim (-M\beta t, -(M/E)\alpha t) \text{ as } t \rightarrow \infty \tag{1.12}$$

and $x(t) - M = O(e^{-M\beta t})$ and $y(t) = O(e^{-(M/E)\alpha t})$.

(iii) When $N = -M/E > 0$, $(x(t), y(t)) \rightarrow (0, N)$ as $t \rightarrow \infty$ and

$$(\log x(t), \log(y(t) - N)) \sim (-EN\beta t, -N\alpha t) \text{ as } t \rightarrow \infty \tag{1.13}$$

and $x(t) = O(e^{-EN\beta t})$ and $y(t) - N = O(e^{-N\alpha t})$.

Remark. When the time dependent coefficients $a(t)$ and $b(t)$ in (1.1) satisfy $a(t)/b(t) = \text{const} > 0$ for $t \geq 0$, we can obtain the similar representation formula of solution $(x(t), y(t))$ of (1.1) replaced αt and βt in (1.6)–(1.10) by $A(t)$ and $B(t)$, respectively.

In [4], Ito, Ogiwara and Usami have derived the following asymptotic forms (1.14) and (1.15) of solution $(x(t), y(t))$ of (1.1) decaying to the origin $(0, 0)$, like (1.11) for (1.1) with constant coefficients :

(i) If $a(t)$ and $b(t)$ satisfy (1.2) and $\lim_{t \rightarrow \infty} a(t)/b(t) = \text{const} > 0$, then

$$(x(t), y(t)) \sim (B(t)^{-1}, A(t)^{-1}) \text{ as } t \rightarrow \infty. \tag{1.14}$$

(ii) If $a(t)$ and $b(t)$ are of class C^1 and satisfy

$$\left(\frac{a(t)}{b(t)}\right)' \leq 0 \text{ for large } t$$

and

$$\lim_{t \rightarrow \infty} \frac{a(t)B(t)}{A(t)b(t)} = k > 0 \text{ and } \lim_{t \rightarrow \infty} \left(\frac{a(t)B(t)}{A(t)b(t)}\right)' \frac{B(t)}{b(t)} = 0,$$

then

$$(x(t), y(t)) \sim (kB(t)^{-1}, k^{-1}A(t)^{-1}) \quad \text{as } t \rightarrow \infty. \quad (1.15)$$

However, there is no known research work related to asymptotic forms of solutions of (1.1) decaying to a point other than the origin $(0, 0)$, like (1.12) and (1.13) for (1.4) with constant coefficients.

The notations we use in this paper are standard. Positive constants will be denoted by C and will change from line to line.

2 Results

We will give asymptotic forms of solutions of (1.1) decaying to a point on the x -axis and y -axis.

Theorem 2.1 *Let E, M and N be constants. Assume that $a(t)$ and $b(t)$ satisfy (1.2) and*

$$\lim_{t \rightarrow \infty} \frac{a(t)}{b(t)} = E > 0. \quad (2.1)$$

Then, we have the following :

(i) *For $M > 0$, every solution $(x(t), y(t))$ of (1.1) decaying to $(M, 0)$ has the asymptotic form*

$$(\log(x(t) - M), \log y(t)) \sim (-MB(t), -(M/E)A(t)) \quad \text{as } t \rightarrow \infty. \quad (2.2)$$

(ii) *For $N > 0$, every solution $(x(t), y(t))$ of (1.1) decaying to $(0, N)$ has the asymptotic form*

$$(\log x(t), \log(y(t) - N)) \sim (-ENB(t), -NA(t)) \quad \text{as } t \rightarrow \infty. \quad (2.3)$$

Proof. (i) Let $M > 0$ and $(x(t), y(t)) \rightarrow (M, 0)$ as $t \rightarrow \infty$. By L'Hospital's rule, we have from (2.1) that

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{x(t) - M}{y(t)} &= \lim_{t \rightarrow \infty} \frac{x'(t)}{y'(t)} = \lim_{t \rightarrow \infty} \frac{-a(t)x(t)y(t)}{-b(t)x(t)y(t)} \\ &= \lim_{t \rightarrow \infty} \frac{a(t)}{b(t)} = E, \end{aligned} \quad (2.4)$$

and hence, we obtain from (2.1) and (2.4) that

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\log(x(t) - M)}{-MB(t)} &= \lim_{t \rightarrow \infty} \frac{(x(t) - M)^{-1}x'(t)}{-Mb(t)} \\ &= \lim_{t \rightarrow \infty} \frac{(x(t) - M)^{-1}(-a(t)x(t)y(t))}{-Mb(t)} \\ &= \lim_{t \rightarrow \infty} \frac{a(t)}{b(t)} \frac{y(t)}{x(t) - M} \frac{x(t)}{M} = 1, \end{aligned}$$

which implies $\log(x(t) - M) \sim -MB(t)$ as $t \rightarrow \infty$.

On the other hand, by L'Hospital's rule again, we have from (2.1) that

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\log y(t)}{-(M/E)A(t)} &= \lim_{t \rightarrow \infty} \frac{y(t)^{-1}y'(t)}{-(M/E)a(t)} = \lim_{t \rightarrow \infty} \frac{y(t)^{-1}(-b(t)x(t)y(t))}{-(M/E)a(t)} \\ &= E \lim_{t \rightarrow \infty} \frac{b(t)}{a(t)} \frac{x(t)}{M} = 1, \end{aligned}$$

which implies $\log y(t) \sim -(M/E)A(t)$ as $t \rightarrow \infty$.

(ii) Next, let $N > 0$ and $(x(t), y(t)) \rightarrow (0, N)$ as $t \rightarrow \infty$. By L'Hospital's rule, we have from (2.1) that

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\log x(t)}{-ENB(t)} &= \lim_{t \rightarrow \infty} \frac{x(t)^{-1}x'(t)}{-ENb(t)} = \lim_{t \rightarrow \infty} \frac{x(t)^{-1}(-a(t)x(t)y(t))}{-ENb(t)} \\ &= \frac{1}{E} \lim_{t \rightarrow \infty} \frac{a(t)}{b(t)} \frac{y(t)}{N} = 1, \end{aligned}$$

which implies $\log x(t) \sim -ENB(t)$ as $t \rightarrow \infty$.

On the other hand, by L'Hospital's rule again, we have from (2.1) that

$$\lim_{t \rightarrow \infty} \frac{x(t)}{y(t) - N} = \lim_{t \rightarrow \infty} \frac{x'(t)}{y'(t)} = \lim_{t \rightarrow \infty} \frac{a(t)}{b(t)} = E, \quad (2.5)$$

and hence, we obtain from (2.1) and (2.5) that

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\log(y(t) - N)}{-NA(t)} &= \lim_{t \rightarrow \infty} \frac{(y(t) - N)^{-1}y'(t)}{-Na(t)} \\ &= \lim_{t \rightarrow \infty} \frac{(y(t) - N)^{-1}(-b(t)x(t)y(t))}{-Na(t)} \\ &= \lim_{t \rightarrow \infty} \frac{b(t)}{a(t)} \frac{x(t)}{y(t) - N} \frac{y(t)}{N} = 1, \end{aligned}$$

which implies $\log(y(t) - N) \sim -NA(t)$ as $t \rightarrow \infty$. \square

Theorem 2.2 *Let M be a constant. Assume that $a(t)$ and $b(t)$ are of class C^1 and satisfy (1.2) and*

$$\left(\frac{a(t)}{b(t)} \right)' \leq 0 \quad \text{for large } t. \quad (2.6)$$

Then, for $M > 0$, every solution $(x(t), y(t))$ of (1.1) decaying to $(M, 0)$ has

$$x(t) - M = O(e^{-MB(t)}). \quad (2.7)$$

In addition, if there exists a positive constant K such that

$$\lim_{t \rightarrow \infty} \frac{a(t)^2 e^{MB(t)}}{b(t)^2 e^{KA(t)}} = \text{const} > 0, \quad (2.8)$$

then

$$y(t) = O(e^{-KA(t)}). \quad (2.9)$$

Proof. Since $x(t) \rightarrow M$ as $t \rightarrow \infty$ and $(-x(t))' = (a(t)/b(t))(-y(t))'$, it follows that

$$\begin{aligned} x(t) - M &= \int_t^\infty (-x(s))' ds = \int_t^\infty \frac{a(s)}{b(s)} (-y(s))' ds \\ &= \frac{a(t)}{b(t)} y(t) + \int_t^\infty \left(\frac{a(s)}{b(s)} \right)' y(s) ds \end{aligned}$$

for large t , and from (2.6) that there exists $t_1 > 0$ such that

$$y(t) \geq \frac{b(t)}{a(t)} (x(t) - M) \quad \text{for } t \geq t_1.$$

Then we have

$$x'(t) = -a(t)x(t)y(t) \leq b(t)x(t)(x(t) - M)$$

for $t \geq t_1$. Solving this differential inequality of a separable type on $[t_1, t]$, we obtain

$$\frac{1}{M} \log \frac{x(t)}{x_1} \frac{x_1 - M}{x(t) - M} \geq B(t) - B_1$$

and

$$x(t) \leq \frac{M}{1 - (1 - M/x_1)e^{-M(B(t) - B_1)}} \leq Ce^{-MB(t)},$$

where we use symbols $x_1 = x(t_1)$ and $B_1 = B(t_1)$, and hence,

$$x(t) - M \leq \frac{(1 - M/x_1)e^{-M(B(t) - B_1)}}{1 - (1 - M/x_1)e^{-M(B(t) - B_1)}} \leq Ce^{-MB(t)} \quad (2.10)$$

for $t \geq t_1$, which implies (2.7).

On the other hand, since $y(t) \rightarrow 0$ as $t \rightarrow \infty$ and $(-y(t))' = (b(t)/a(t)) \cdot (-x(t) - M)'$, it follows that

$$\begin{aligned} y(t) &= \int_t^\infty (-y(s))' ds = \int_t^\infty \frac{b(s)}{a(s)} (-x(s) - M)' ds \\ &= \frac{b(t)}{a(t)} (x(t) - M) + \int_t^\infty \left(\frac{b(s)}{a(s)} \right)' (x(s) - M) ds \end{aligned} \quad (2.11)$$

for large t . Here, since it follows from (2.8) that

$$0 \leq \frac{b(t)}{a(t)} e^{-MB(t)} = \frac{a(t)}{b(t)} \frac{b(t)^2}{a(t)^2} e^{-MB(t)} \leq C e^{-KA(t)} \quad \text{for large } t, \quad (2.12)$$

and from $(b(t)/a(t))' = -(b(t)/a(t))^2 (a(t)/b(t))' \geq 0$ for large t and (2.8) that

$$\begin{aligned} 0 &\leq \int_t^\infty \left(\frac{b(s)}{a(s)} \right)' e^{-MB(s)} ds \\ &= -\frac{b(t)}{a(t)} e^{-MB(t)} + \int_t^\infty \frac{b(s)}{a(s)} Mb(s) e^{-MB(s)} ds \\ &\leq -\frac{b(t)}{a(t)} e^{-MB(t)} + C \int_t^\infty Ka(s) e^{-KA(s)} ds \\ &= -\frac{b(t)}{a(t)} e^{-MB(t)} + C \int_t^\infty (-e^{-KA(s)})' ds \\ &\leq C e^{-KA(t)} \quad \text{for large } t, \end{aligned} \quad (2.13)$$

we obtain from (2.10)–(2.13) that

$$y(t) \leq C e^{-KA(t)} + C e^{-KA(t)} \quad \text{for large } t,$$

which implies (2.9). \square

Remark. (i) When $a(t) = \alpha > 0$ and $b(t) = \beta > 0$, we see that $(a(t)/b(t))' = 0$ and the limit value of (2.8) is $\alpha^2/\beta^2 > 0$ by taking $K = M\beta/\alpha$.

(ii) When $a(t) = (1+t)^{-1}$ and $b(t) = (e+t)^{-1}$, we see that $(a(t)/b(t))' < 0$ and the limit value of (2.8) is $e^{-M} > 0$ by taking $K = M$.

By the similar argument of Theorem 2.2 we have the following theorem.

Theorem 2.3 *Let N be a constant. Assume that $a(t)$ and $b(t)$ are of class C^1 and satisfy (1.2) and*

$$\left(\frac{b(t)}{a(t)} \right)' \leq 0 \quad \text{for large } t.$$

Then, for $N > 0$, every solution $(x(t), y(t))$ of (1.1) decaying to $(0, N)$ has

$$y(t) - N = O(e^{-NA(t)}).$$

In addition, if there exists a positive constant K such that

$$\lim_{t \rightarrow \infty} \frac{b(t)^2 e^{NA(t)}}{a(t)^2 e^{KB(t)}} = \text{const} > 0,$$

then

$$x(t) = O(e^{-KB(t)}).$$

References

- [1] M.J. Artelli and R.F. Deckro, Modeling the Lanchester laws with system dynamics, *J. Defence Modeling and Simulation*, **5** (2008) 1–20.
- [2] D.R. Howes and R.M. Thrall, A theory of ideal linear weights for heterogeneous combat forces, *Naval Research Logistics Quarterly*, **20** (1973) 645–659.
- [3] G. Isac and A. Gosselin, A military application of viability: winning cones, differential inclusions, and Lanchester type models for combats, *Pareto Optimality, Game Theory and Equilibria*, 759–797, Springer Optim. Appl., 17, Springer, New York 2008.
- [4] T. Ito, T.Ogiwara, and H. Usami, Asymptotic properties of solutions of a Lanchester-type model, *Differential Equations & Applications* **12** (2020) 1–12.
- [5] F.W. Lanchester, *Aircraft in Warfare: The Dawn of the Fourth Arm*, 1916 (Reprinted by Forgotten Books, 2015).
- [6] K.Y. Lin and N. J. MacKay, The optimal policy for the one-against-many heterogeneous Lanchester, *Oper. Res. Lett.* **42** (2014) 473–477.
- [7] N.J. MacKay, Lanchester models for mixed forces with semi-dynamical target allocation, *J. Oper. Res. Soc.* **60** (2009) 1421–1427.
- [8] T. Ogiwara and H. Usami, On the behavior of solutions for Lanchester square-law models with time-dependent coefficients, *Josai Mathematical Monographs*, **11** (2018) 15–25.
- [9] J.G. Taylor, *Lanchester Models of Warfare, I and II*, Operations Research Society of America 1983.
- [10] J.G. Taylor, Battle-outcome prediction for an extended system of Lanchester type differential equations, *J. Math. Anal. Appl.*, **103** (1984) 371–379.
- [11] H.T. Tran and H. Usami, Asymptotic behavior of positive solutions of a Lanchester-type model, *Differential Equations and Applications*, **9** (2017) 241–252.